

RATIONALLY SMOOTH ELEMENTS OF COXETER GROUPS AND TRIANGLE GROUP AVOIDANCE

EDWARD RICHMOND AND WILLIAM SLOFSTRA

ABSTRACT. We study a family of infinite-type Coxeter groups defined by the avoidance of certain rank 3 parabolic subgroups. For this family, rationally smooth elements can be detected by looking at only a few coefficients of the Poincaré polynomial. We also prove a factorization theorem for the Poincaré polynomial of rationally smooth elements. As an application, we show that a large class of infinite-type Coxeter groups have only finitely many rationally smooth elements. Explicit enumerations and descriptions of these elements are given in special cases.

1. INTRODUCTION

Let W be a Coxeter group with finite generating reflection set S , and let ℓ and $\leq$ denote the length function and Bruhat order on W , respectively. Let $e \in W$ denote the identity of W . By definition, W is the group generated by S satisfying relations $(st)^{m_{st}} = e$, where $m_{st} \in \{1, 2, 3, \dots, \infty\}$ such that $m_{st} = 1$ if and only if $s = t$. If $m_{st} = \infty$, then by convention the relation $(st)^\infty = e$ is omitted. The Poincaré series

$$P_w(q) = \sum_{x \leq w} q^{\ell(x)}$$

of an element $w \in W$ is a polynomial of degree $\ell(w)$. An element w is said to be *palindromic* (or *rationally smooth*) if the coefficients of $P_w(q)$ are the same whether read from top degree to bottom degree, or in reverse.¹ In other words, if we write $P_w(q) = \sum a_i q^i$, then w is palindromic when $a_i = a_{\ell(w)-i}$ for all i .

An important question in the combinatorics of Coxeter groups is to describe the set of palindromic elements of W . This question stems from its connection with the topology of Schubert varieties. A Coxeter group is *crystallographic* if $m_{st} \in \{2, 3, 4, 6, \infty\}$ for all $s \neq t$. If W is crystallographic, then it can be realized as the Weyl group of a Kac-Moody algebra. The Schubert subvarieties of the full flag variety corresponding to this algebra are indexed by the elements of W . Carrell and Peterson prove that the Schubert variety indexed by w is rationally smooth if and only if w is palindromic [9]. Furthermore, w is palindromic if and only if the Kazhdan-Lusztig polynomial indexed by (x, w) is equal to 1 for all $x \leq w$ [11, 12]. If W is crystallographic, then it is sufficient that the Kazhdan-Lusztig polynomial indexed by (e, w) be equal to 1 [9]. For Schubert varieties of simply laced types A , D and E , the notion of smooth and rationally smooth are equivalent. For finite Weyl groups, the palindromic elements are well understood. In particular, they can be

¹The term rationally smooth seems to be more common in the literature; we use the term palindromic to be inclusive of the non-crystallographic case.

characterized using permutation pattern-avoidance in classical types A, B, C , and D and using root system avoidance in all types [3, 4, 5, 13]. The characterization using permutation pattern avoidance has recently been extended to the affine type A case as well [2]. The generating series for the number of palindromic elements in A_n , as n varies, is also known [3, 8, 16].

While the theory of palindromic elements is well-developed for finite and affine Coxeter groups, the situation for general Coxeter groups is quite different. In particular, it seems to be quite difficult to determine whether or not an element of a general Coxeter group is rationally smooth. In this paper, we introduce a family of Coxeter groups (mostly) outside the finite and affine cases, for which it is possible to determine if an element is rationally smooth by looking at just a few coefficients of the Poincaré polynomial. The family in question is defined as the set of all Coxeter groups which do not contain certain triangle groups as standard parabolic subgroups. A *triangle group* is a Coxeter group with $|S| = 3$. Triangle groups arise naturally in arithmetic geometry and the study of tessellations of triangles on Riemann surfaces, see e.g. [1]. We will denote a triangle group by the triple (m_{rs}, m_{rt}, m_{st}) where $S = \{r, s, t\}$. We say a Coxeter group W *contains the triangle* (a, b, c) if there exists a subset $\{r, s, t\} \subseteq S$ such that $(a, b, c) = (m_{rs}, m_{rt}, m_{st})$. If S contains no such subset, then we say W *avoids the triangle* (a, b, c) . We are interested in the groups which avoid the following special set of triangle groups:

$$\text{HQ} := \{(2, b, c) \mid b, c \geq 3 \text{ and } b < \infty\}$$

The set HQ (Hecke quotients) is the set of quotients of the Hecke triangle group $(2, p, \infty)$, $p \geq 3$, which is a generalization of the well-known modular group $(2, 3, \infty)$. Every finite Coxeter group of rank ≥ 3 contains a triangle in HQ, and the same is true of affine Coxeter groups, with the exception of $(3, 3, 3)$, which is the affine group $\hat{A}_2$. However, there are many crystallographic Coxeter groups which do avoid HQ; for example, any Coxeter group with no commuting relations (i.e. $m_{st} \geq 3$ for all $s \neq t$) avoids HQ. Any Coxeter group defined by only by commuting and infinite relations also avoids HQ.

To state our main theorem, we make the following definition:

Definition 1.1. *Let w be an element of a Coxeter group W , and write $P_w(q) = \sum a_i q^i$ for the Poincaré polynomial of w . We say that w is k -palindromic if $a_i = a_{\ell(w)-i}$ for all $0 \leq i < k$.*

Note that if $k = \infty$, then we recover the usual notion of palindromic elements, and that every element is 1-palindromic with $a_0 = a_{\ell(w)} = 1$. If W is crystallographic, then k -palindromicity can be detected from the Kazhdan-Lusztig polynomial. Let $T_{e,w} = 1 + \sum_{i \geq 0} b_i q^i$ be the Kazhdan-Lusztig polynomial indexed by (e, w) . A theorem of Bjorner and Ekedahl states that, for crystallographic groups, an element $w \in W$ is k -palindromic if and only if $b_i = 0$ for $0 \leq i < k$ [7] (note that $b_0 = 0$ always).

We now state the main theorem:

Theorem 1.2. *Let W be a Coxeter group which avoids all triangle groups in HQ. Then every 4-palindromic $w \in W$ is palindromic.*

Furthermore, if W avoids all triangle groups $(3, 3, c)$ where $3 < c < \infty$, then every 2-palindromic $w \in W$ is palindromic.

Given a Coxeter group, it is natural to ask whether there is a number k such that every k -palindromic element is palindromic. This question appears to be open in general. Billey and Postnikov have conjectured that if W is a finite simply-laced Weyl group with n generators, then every $(n + 1)$ -palindromic element of W is palindromic [4]. In type A_n , it is known that every $(n - 1)$ -palindromic element is palindromic [4].

The proof of Theorem 1.2 is based on a factorization theorem for the Poincaré polynomial of 2-palindromic elements in Coxeter groups which avoid HQ. In the classical groups of finite type A, B, C , and D , it is known that the Poincaré polynomial of a rationally smooth element factors into a product of q -integers (see Equation (2)) [5, 10]. In fact, it is possible to see this factorization combinatorially, writing each palindromic element w as a reduced product $w_1 \cdots w_{|S|}$, such that each q -integer factor of the Poincaré polynomial equals the (relative) Poincaré polynomial of the w_i 's. We prove a similar result for 2-palindromic elements in Coxeter groups which avoid HQ. This result has a number of applications. For example, we show there are many infinitely large Coxeter groups with only a finite number of palindromic elements. We also give explicit descriptions of palindromic elements in special cases. In the case of uniform Coxeter groups $W(m, n)$, defined by $m_{st} = m$ for all $s \neq t$ and $|S| = n$, we calculate the generating series for the number of palindromic elements weighted by length. Formulas for these generating series are stated in Propositions 3.8 and 3.9. We also observe that the HQ-avoiding groups form the largest class of Coxeter groups for which our factorization theorem can hold.

1.1. Organization. Section 2 contains some background material and elementary lemmas used to state the factorization theorem. Section 3 states the main factorization theorem and its consequences, including the proof of Theorem 1.2 and enumerative results. In Section 4, we consider triangle groups in the set HQ and prove the main results cannot hold for any Coxeter group containing these triangle groups. Section 5 gives some elementary lemmas on the descent sets of Coxeter groups avoiding HQ. Finally, Section 6 proves the main factorization theorem.

2. BACKGROUND AND TERMINOLOGY

Let W be a Coxeter group with simple generator set S . For basic facts on Coxeter groups, we refer the reader to [6]. Let $\ell(w)$ denote the length of $w \in W$. We say $w = uv \in W$ is a *reduced factorization* if $\ell(w) = \ell(u) + \ell(v)$. A special type of reduced factorization can be constructed from any subset $J \subseteq S$. Let W_J denote the standard parabolic subgroup of W generated by J . Let W^J denote the set of minimal length coset representatives of $W_J \backslash W$. Every element $w \in W$ can be written uniquely as $w = uv$ where $u \in W_J$, $v \in W^J$ and $\ell(w) = \ell(u) + \ell(v)$. We call this reduced factorization of w the *parabolic decomposition* with respect to J .

Let $\leq$ denote the Bruhat order on W . If $u \leq v \in W$, then the interval $[u, v]$ denotes the set of elements $x \in W$ such that $u \leq x \leq v$. For any $w \in W$ we can define the Poincaré polynomial

$$P_w(q) := \sum_{x \in [e, w]} q^{\ell(x)}.$$

The Poincaré polynomial relative to $J \subseteq S$ of an element $w \in W$ is defined to be

$$P_w^J(q) := \sum_{x \in [e, w] \cap W^J} q^{\ell(x)}.$$

If $w \in W^J$, then $P_w^J(q)$ is a polynomial of degree $\ell(w)$. If $J = \emptyset$, then $P_w^J(q) = P_w(q)$. Recall that for any J , the poset $[e, w] \cap W_J$ has a unique maximal element. The following proposition is due to Billey and Postnikov in [4, Theorem 6.4].

Proposition 2.1 ([4]). *Let $J \subseteq S$ and let $w = uv$ be a parabolic decomposition with respect to J . Then u is the unique maximal element of $[e, w] \cap W_J$ if and only if*

$$P_w(q) = P_u(q) \cdot P_v^J(q).$$

While the proof of Proposition 2.1 given in [4] is stated only for finite Weyl groups, it easily extends to all Coxeter groups. A parabolic decomposition $w = uv$ is called a *BP-decomposition* of w if u is the unique maximal element of $[e, w] \cap W_J$.

For any $w \in W$, define the sets

$$\begin{aligned} \mathbf{S}(w) &:= \{u \leq w \mid \ell(u) = 1\} \\ \mathbf{D}(w) &:= \{u \leq w \mid \ell(u) = \ell(w) - 1\} \\ D_R(w) &:= \{s \in S \mid \ell(ws) < \ell(w)\} \\ D_L(w) &:= \{s \in S \mid \ell(sw) < \ell(w)\}. \end{aligned}$$

The sets $\mathbf{S}(w)$ and $\mathbf{D}(w)$ are known as the support and divisor sets of w . The sets $D_R(w)$ and $D_L(w)$ are called the *right* and *left descent sets* of w respectively and are contained in $\mathbf{S}(w)$. We use these sets to give an equivalent characterization of a BP-decomposition.

Lemma 2.2. *A parabolic decomposition $w = uv$ is a BP-decomposition if and only if $\mathbf{S}(v) \cap J \subseteq D_R(u)$.*

Proof. If $w = uv$ is a BP-decomposition, then u is the unique longest element of $[e, w] \cap W_J$. If there exists $x \in \mathbf{S}(v) \cap J$ and $x \notin D_R(u)$, then $\ell(ux) = \ell(u) + 1$ and $ux \in [e, w] \cap W_J$ which is a contradiction.

Conversely, assume that $\mathbf{S}(v) \cap J \subseteq D_R(u)$ and let $\bar{u}$ denote the maximal element in $[e, w] \cap W_J$. Since $\bar{u}$ is unique, we have that $u \leq \bar{u}$. We now show that $\bar{u} \leq u$. Let

$$\bar{u} = u'v'$$

be a reduced factorization which maximizes $\ell(u')$ under the conditions that $u' \leq u$ and $v' \leq v$. Suppose that $v' \neq e$. Then there exists $y \in D_L(v') \setminus D_R(u')$. By assumption, we have that $y \in D_R(u)$. Taking a reduced decomposition for u with

y appearing at the end, we see that $u' \leq uy$, and hence u' can be extended, a contradiction. $\square$

We remark that one direction of Lemma 2.2 is proved in [15, Lemma 10]. Another property of BP-decompositions is the following lemma.

Lemma 2.3. *Let $J_1 \subseteq J_2 \subseteq S$ and let $v_1v_2v_3$ be a reduced factorization such that v_1v_2 and $(v_1v_2)(v_3)$ are BP-decompositions with respect to J_1 and J_2 respectively. Then $v_1(v_2v_3)$ is a BP-decomposition with respect to J_1 .*

Proof. By definition, we have v_1v_2 is maximal in $[e, v_1v_2v_3] \cap W_{J_2}$. In particular, if u denotes the maximal element in $[e, v_1v_2v_3] \cap W_{J_1}$, then $u \leq v_1v_2$ since $W_{J_1} \subseteq W_{J_2}$. But now u is maximal in $[e, v_1v_2] \cap W_{J_1}$, which implies that $u = v_1$. $\square$

Clearly, if $P_w(q) = \sum a_i q^i$, then $|\mathbf{S}(w)| = a_1$ and $|\mathbf{D}(w)| = a_{\ell(w)-1}$. We now consider a special class of parabolic decompositions.

Definition 2.4. *We say that $w = uv$, a parabolic decomposition with respect to J , is a Grassmannian factorization if $J = \mathbf{S}(u)$ and $|\mathbf{S}(w)| = |\mathbf{S}(u)| + 1$*

It is easy to see that every element $w \in W$ of length ≥ 2 has a Grassmannian factorization. The term ‘‘Grassmannian’’ comes from the fact that v is a Grassmannian element of W which, by definition, has $|D_L(v)| = 1$. Note that a Grassmannian factorization is not necessarily a BP-decomposition. Although elementary, this concept is quite useful. For example, we can use it to prove:

Lemma 2.5. $|\mathbf{D}(w)| \geq |\mathbf{S}(w)|$.

Proof. We proceed by induction on $\ell(w)$. The proposition is true if $\ell(w) = 1$, so suppose $\ell(w) \geq 2$. Let $w = uv$ be a Grassmannian factorization with respect to J . By induction, $|\mathbf{D}(u)| \geq |\mathbf{S}(u)|$.

If $u' \in \mathbf{D}(u)$, then $u'v \in \mathbf{D}(w)$, since $v \in W^J$. Now v is not the identity, so we can write $v' = vs \in W^J$ with $s \in S$ and $\ell(v') = \ell(v) - 1$. Consequently $uv' \in \mathbf{D}(w)$. Moreover, $uv' \neq u'v$ for any $u' \in \mathbf{D}(u)$ since they are both parabolic decompositions with respect to J and $u \neq u'$. Hence

$$(1) \quad |\mathbf{D}(w)| \geq |\mathbf{D}(u)| + 1 \geq |\mathbf{S}(u)| + 1 = |\mathbf{S}(w)|.$$

This completes the proof. $\square$

We remark that, for crystallographic Coxeter groups, Bjorner and Ekedahl prove a much stronger version of Lemma 2.5 concerning all the coefficients of $P_w(q)$ [7, Theorem A].

We can continue to decompose any Grassmannian factorization $w = uv$ by taking a Grassmannian factorization of u . We say that

$$w = v_1v_2 \cdots v_{|\mathbf{S}(w)|}$$

is a *complete Grassmannian factorization* of w if for every $i < |\mathbf{S}(w)|$, we have that $(v_1 \cdots v_i)(v_{i+1})$ is a Grassmannian factorization. Observe that if each $(v_1 \cdots v_i)(v_{i+1})$

is also a BP-decomposition, then by Lemma 2.3, we have $(v_1 \cdots v_i)(v_{i+1} \cdots v_k)$ is BP decomposition for any $i < k \leq |\mathbf{S}(w)|$.

By definition, w is 2-palindromic if and only if $|\mathbf{D}(w)| = |\mathbf{S}(w)|$. The following lemma gives an inductive characterization of the 2-palindromic property.

Lemma 2.6. *Suppose that $w = uv$ is a Grassmannian factorization. Then w is 2-palindromic if and only if u is 2-palindromic and $|u \cdot \mathbf{D}(v) \cap \mathbf{D}(w)| = 1$.*

Proof. Equality holds in Equation (1) if and only if $|\mathbf{D}(u)| = |\mathbf{S}(u)|$ and $u \cdot \mathbf{D}(v) \cap \mathbf{D}(w) = \{uvs\}$ where $s \in D_R(v)$. $\square$

3. THE FACTORIZATION THEOREM

The main technical theorem of this paper is the following:

Theorem 3.1. *Suppose that W avoids all triangle groups in HQ. Let $w \in W$ be 2-palindromic and fix a Grassmannian factorization $w = uv$ with respect to $J \subseteq S$. Then $w = uv$ is a BP-decomposition with respect to J such that $|\mathbf{S}(v)| \leq 3$.*

Moreover, if $|\mathbf{S}(v)| = 3$ and $\mathbf{S}(v) = \{r, s, t\}$, then one of the following is true:

- (1) $v = trv'$ with $v' = \underbrace{stst \dots}_{m_{st}-1}$ where $\mathbf{S}(v)$ generates the triangle group $(3, m_{rs}, m_{st})$ with $m_{rt} = 3$ and $3 \leq m_{st} < \infty$, $3 \leq m_{rs} \leq \infty$.
- (2) $v = rstrv'$ with $v' = \underbrace{stst \dots}_{m_{st}-1}$ where $\mathbf{S}(v)$ generates the triangle group $(3, 3, m_{st})$ with $3 < m_{st} < \infty$.
- (3) $v = strstr \dots$ is a spiral word² of even length where $\mathbf{S}(v)$ generates the triangle group $(3, 3, 3)$.

Theorem 3.1 says that if W avoids triangle groups in HQ, then the Poincaré polynomial $P_w(q)$ of a 2-palindromic element $w \in W$ factors along any Grassmannian factorization of $w = uv$. Moreover, the possibilities for the factor $P_v^J(q)$ is limited by the fact that $|\mathbf{S}(v)| \leq 3$. Note that parts (1) and (3) of the theorem overlap when $m_{rs} = m_{st} = 3$. The proof of this theorem is the focus of Section 6. The remainder of this section is devoted to consequences of Theorem 3.1.

Fix a 2-palindromic element $w \in W$ and a Grassmannian factorization $w = uv$ with respect to $J \subseteq S$. Theorem 3.1 can be used, together with Lemma 2.6, to completely determine the polynomial $P_w(q)$. By Theorem 3.1 and Proposition 2.1, we have that

$$P_w(q) = P_u(q) \cdot P_v^J(q),$$

so it suffices to characterize all possible polynomials $P_v^J(q)$. For any integer $k \geq 1$ define the q -integer

$$(2) \quad [k]_q := 1 + q + \dots + q^{k-1}.$$

If $|\mathbf{S}(v)| \leq 2$, then any $v' \leq v$ where $v' \in W^J$ is given by a prefix of the unique reduced word of v . This implies

$$(3) \quad P_v^J(q) = [\ell(v) + 1]_q.$$

²A spiral word is word which cycles through a set of generators in a fixed order.

If $|\mathbf{S}(v)| = 3$, it suffices to compute $P_v^J(q)$ in all the cases of Theorem 3.1. We have the following lemma.

Lemma 3.2. *Suppose we have $w = uv$ as in Theorem 3.1 with $|\mathbf{S}(v)| = 3$. Then the following are true:*

- (1) *If v satisfies the conditions in Theorem 3.1 part (1), then*

$$P_v^J(q) = [\ell(v) + 1]_q + q^2[\ell(v) - 3]_q.$$

- (2) *If v satisfies the conditions in Theorem 3.1 part (2), then*

$$P_v^J(q) = [\ell(v) + 1]_q + q^2[\ell(v) - 3]_q + q^4[\ell(v) - 6]_q.$$

- (3) *If v satisfies the conditions in Theorem 3.1 part (3) with $k = \left\lfloor \frac{\ell(v)}{4} \right\rfloor$, then*

$$P_v^J(q) = \sum_{i=0}^k q^{2i} [\ell(v) - 4i + 1]_q.$$

Proof. Part (3) is proved in [14, Proposition 2.4] where certain Poincaré polynomials of Schubert varieties in the affine Grassmannian of type A are calculated. Parts (1) and (2) can be deduced from elementary counting arguments of the sets

$$\{v' \in W^J \cap [e, v] \mid \ell(v') = i\}.$$

In particular, for part (1), there are two q -integer contributions from reduced subwords of the form

$$tr \underbrace{stst \dots}_k \quad \text{and} \quad \underbrace{tstst \dots}_k.$$

For part (2) there are three q -integer contributions from reduced subwords of the form

$$r \underbrace{tsts \dots}_k \quad \text{and} \quad r \underbrace{stst \dots}_k \quad \text{and} \quad rstr \underbrace{stst \dots}_k.$$

□

The polynomials in parts (1) and (3) of the lemma are palindromic, while the polynomial in part (2) is 3-palindromic but not 4-palindromic. We now prove the theorem stated in the introduction.

Proof of Theorem 1.2. Suppose that W avoids all triangles in HQ. Let $w = v_1 v_2 \dots v_{|\mathbf{S}(w)|} \in W$ be a complete Grassmannian factorization. Then by Theorem 3.1 and Proposition 2.1, we have that

$$P_w(q) = \prod_{i=1}^{|\mathbf{S}(w)|} P_{v_i}^{J_i}(q).$$

where $J_i := \mathbf{S}(v_1) \cup \dots \cup \mathbf{S}(v_{i-1})$ and $J_1 := \emptyset$. Moreover, the factors $P_{v_i}^{J_i}(q)$ are given by either Equation (3) or by parts (1)–(3) of Lemma 3.2. Now $P_w(q)$ is 4-palindromic if the polynomial in Lemma 3.2 part (2) does not appear as one of the factors $P_{v_i}^{J_i}(q)$. Since all other possible choices for $P_{v_i}^{J_i}(q)$ are palindromic, we have that $P_w(q)$ is 4-palindromic if and only if it is palindromic. This proves part (1) of Theorem 1.2.

If W also avoids the triangles of the form $(3, 3, c)$, then Lemma 3.2 part (2) is never an option for $P_{v_i}^{J_i}(q)$. Hence every 2-palindromic $w \in W$ is palindromic. This completes the proof. $\square$

3.1. Examples. Consider the Coxeter group W with $S = \{s_1, s_2, s_3, s_4\}$ defined by the Dynkin diagram in Figure 1.




FIGURE 1.

Unlabeled edges are assumed to have label $m_{st} = 3$ and if there is no edge between s and t , then $m_{st} = 2$. Clearly, W avoids all triangle groups in HQ and hence we can apply Theorem 3.1 to compute Poincaré polynomials.

Example 3.3. Let $w = s_1 s_2 s_1 s_3 s_2 s_1 s_3 s_2 s_1 s_4$. Then w is 2-palindromic with $|\mathbf{S}(w)| = |\mathbf{D}(w)| = 4$. The following is a complete Grassmannian factorization:

$$w = \underbrace{(s_1)}_{v_1} \underbrace{(s_2 s_1)}_{v_2} \underbrace{(s_3 s_2 s_1 s_3 s_2 s_1)}_{v_3} \underbrace{(s_4)}_{v_4}.$$

The corresponding Poincaré polynomial factorization is

$$\begin{aligned} P_w(q) &= [2]_q [3]_q ([7]_q + q^2 [3]_q) [2]_q \\ &= (1+q)(1+q+q^2)(1+q+2q^2+2q^3+2q^4+q^5+q^6)(1+q), \end{aligned}$$

so $P_w(q)$ is palindromic.

Example 3.4. Let $w = s_2 s_4 s_2 s_4 s_1 s_2 s_4 s_1 s_2 s_4 s_2$. Then w is 2-palindromic with $|\mathbf{S}(w)| = |\mathbf{D}(w)| = 3$. A complete Grassmannian factorization of w is

$$w = \underbrace{(s_2)}_{v_1} \underbrace{(s_4 s_2 s_4)}_{v_2} \underbrace{(s_1 s_2 s_4 s_1 s_2 s_4 s_2)}_{v_3}.$$

The corresponding Poincaré polynomial factorization is

$$\begin{aligned} P_w(q) &= [2]_q [4]_q ([8]_q + q^2 [4]_q + q^4 [1]_q) \\ &= (1+q)(1+q+q^2+q^3)(1+q+2q^2+2q^3+3q^4+2q^5+q^6+q^7) \end{aligned}$$

Note that $\{s_1, s_2, s_4\}$ generates the triangle group $(3, 3, 4)$. Since $v_3 = s_1 s_2 s_4 s_1 s_2 s_4 s_2$, we have that w is 3-palindromic but not 4-palindromic.

An example of a HQ-avoiding Coxeter group W with commuting relations is given by the Dynkin diagram in Figure 2 below where $p \geq 3$.

Observe that W also avoids all triangle groups of the form $(3, 3, c)$. Hence every 2-palindromic element is palindromic by Theorem 1.2. Moreover, every palindromic




FIGURE 2.

polynomial factors into a product of q -integers. We also remark that W is indecomposable with respect to products and free products of Coxeter groups.

3.2. Enumeration and description of palindromic elements. Theorem 3.1 gives a description of the set of palindromic (resp. 2-palindromic) elements of any HQ-avoiding Coxeter group. Specifically, the palindromic (resp. 2-palindromic) elements are those with a certain Grassmannian factorization. In this section we provide some applications of this idea. We start by proving a corollary of Theorem 3.1 on the finiteness of the number of palindromic elements for all HQ-avoiding Coxeter groups.

Corollary 3.5. *Let (W, S) be a Coxeter group that avoids all triangle groups in HQ. Then W has a finite number of palindromic elements if and only if $m_{st} < \infty$ for all $s, t \in S$ and W avoids the triangle group $(3, 3, 3)$.*

Proof. Theorem 3.1 part (3) implies that the triangle group $(3, 3, 3)$ contains an infinite number of palindromic elements. Also, if $m_{rs} = \infty$, then $W_{\{r, s\}}$ is infinite and every element is palindromic.

Let m_0 denote the largest value of m_{st} for $s, t \in S$. Suppose that W avoids $(3, 3, 3)$ and $m_0 < \infty$. Let $w \in W$ be palindromic with complete Grassmannian factorization $w = v_1 \cdots v_{|S(w)|}$. By Theorem 3.1, we have that each factor v_i has length at most $m_0 + 3$, so

$$\ell(w) < |S(w)|(m_0 + 3) \leq |S|(m_0 + 3)$$

and hence the number of palindromic elements in W is finite. $\square$

Corollary 3.5 also holds if palindromic is replaced by 2-palindromic.

Note that the Grassmannian factorization of an element provided by Theorem 3.1 is not necessarily unique. When $m_{st} \geq 3$ for all $s \neq t$, we give a modified factorization which does not have this problem. To state the modified factorization we need the following definition.

Definition 3.6. *We say a reduced factorization $w = u_1 u_2 \cdots u_d$ is separable if $S(u_i) \cap S(u_j) = \emptyset$ for all $i \neq j$. If no such non-trivial factorization exists, then we say that w is inseparable.*

Given any complete Grassmannian factorization of a palindromic element $w = v_1 \cdots v_{|S(w)|}$, there is a simple method for constructing a separable factorization. Let $(i_1, \dots, i_d)$ denote the subsequence of integers for which $\ell(v_{i_j}) = 1$. Then $w = u_1 \cdots u_d$ is a separable factorization where

$$u_j := v_{i_j} v_{i_j+1} \cdots v_{i_{j+1}-1}$$

and $i_{d+1} := |\mathbf{S}(w)| + 1$. We remark that $\ell(v_1) = 1$ and hence the sequence $(i_1, \dots, i_d)$ is nonempty. Furthermore, each factor u_j is inseparable. For example, let W be defined by the Dynkin diagram in Figure 1 and $w = s_4 s_2 s_4 s_2 s_3 s_1 s_3$. Then $w = u_1 u_2$ given by

$$(4) \quad w = \underbrace{s_4}_{v_1} \underbrace{s_2 s_4 s_2}_{v_2} \underbrace{s_3}_{v_3} \underbrace{s_1 s_3}_{v_4}$$

is a separable factorization. The following corollary follows from Theorem 3.1.

Corollary 3.7. *Let W be a Coxeter group with $m_{st} \geq 3$ for all $s \neq t$, and let $w \in W$ be palindromic. Then w has a unique separable factorization $w = u_1 \cdots u_d$ where each u_i is inseparable and palindromic. Moreover, any complete Grassmannian factorization $u_i = v_1 \cdots v_{|\mathbf{S}(u_i)|}$ is unique up to choice of v_1 .*

Proof. Any element w has a separable factorization $w = u_1 \cdots u_d$ where each u_i is inseparable. Since $\mathbf{S}(u_i)$ is distinct and W has no commuting braid relations, the factorization is unique. If w is palindromic, then every u_i is palindromic since $(u_1 \cdots u_i)(u_{i+1})$ is a BP-decomposition with respect to $J = S \setminus \mathbf{S}(u_i)$.

Let $u_i = v_1 \cdots v_{|\mathbf{S}(u_i)|}$ be a complete Grassmannian factorization, and let s_j be the unique element of $D_L(v_j)$. Note that $v_1 = s_1$. As mentioned above, since u_i is inseparable, we must have $|\mathbf{S}(v_j)| \geq 2$ for $j = 2, \dots, |\mathbf{S}(u_i)|$. Indeed, if $\mathbf{S}(v_j) = \{s_j\}$ then s_j is the unique right descent of $v_1 \cdots v_j$, since $s_j \notin \mathbf{S}(v_1 \cdots v_{j-1})$. But by Lemma 2.3, $(v_1 \cdots v_j)(v_{j+1} \cdots v_{|\mathbf{S}(u_i)|})$ is a BP decomposition, so

$$\mathbf{S}(v_{j+1} \cdots v_{|\mathbf{S}(u_i)|}) \cap \mathbf{S}(v_1 \cdots v_j) \subset \{s_j\}.$$

Thus $(v_1 \cdots v_{j-1})(v_j \cdots v_{|\mathbf{S}(u_i)|})$ is a separable factorization, which is a contradiction.

We now show that s_j is the unique left descent of $v_j \cdots v_{|\mathbf{S}(u_i)|}$, for $j \geq 2$. Indeed, looking ahead to Lemma 5.3, and using the fact that $|\mathbf{S}(v_j)| \geq 2$, we see that $D_L(v_j \cdots v_{|\mathbf{S}(u_i)|})$ is a subset of $\mathbf{S}(v_j) \setminus \mathbf{S}(v_1 \cdots v_{j-1}) = \{s_j\}$. Hence the sequence $(s_2, \dots, s_{|\mathbf{S}(u_i)|})$ is uniquely determined given the choice of $v_1 = s_1$, and the v_j 's are uniquely determined from the corresponding parabolic decomposition. $\square$

Note that there are at most two complete Grassmannian factorizations of each u_i in Corollary 3.7. For example, taking u_1 in Equation (4), we have

$$u_1 = \underbrace{s_4}_{v_1} \underbrace{s_2 s_4 s_2}_{v_2} = \underbrace{s_2}_{v_1} \underbrace{s_4 s_2 s_4}_{v_2}$$

as the only two complete Grassmannian factorizations.

Corollary 3.7 implies that to count the number of palindromic elements of W , it is sufficient to enumerate elements of W which are inseparable and palindromic. When m_{st} is constant we compute an exponential generating series for the number of palindromic elements. Specifically, let $W(m, n)$ denote the *uniform Coxeter group* such that $|S| = n$ and $m_{st} = m$ for all $s \neq t$. Uniform Coxeter groups satisfy the

property that every 2-palindromic element w is palindromic by Theorem 3.1. Define the generating series

$$\Phi_m(q, t) := \sum_{n, k \geq 0} P_{n, k} \frac{q^k t^n}{n!}$$

where $P_{n, k}$ denotes the number of palindromic $w \in W(m, n)$ of length k . In the case that $m = 2$, we have $W(2, n) \simeq (\mathbb{Z}/2\mathbb{Z})^n$ with every element palindromic, so $P_{n, k} = \binom{n}{k}$. Hence the generating series

$$\Phi_2(q, t) = \exp(qt + t).$$

For $m \geq 3$, define

$$\phi_m(q, t) := \sum_{n, k \geq 1} I_{n, k} \frac{q^k t^n}{n!}$$

where $I_{n, k}$ denotes the number of palindromic $w \in W(m, n)$ of length k that are inseparable with $|\mathcal{S}(w)| = n$. Note that Φ_m and ϕ_m are exponential in t and ordinary in q . Corollary 3.7 implies

Proposition 3.8. *For any $3 \leq m \leq \infty$, the series*

$$\Phi_m(q, t) = \frac{\exp(t)}{1 - \phi_m(q, t)}.$$

The following proposition completes the calculation.

Proposition 3.9. *The exponential generating series for the number of inseparable palindromic elements in $W(m, n)$ is*

$$(5) \quad \phi_m(q, t) = \begin{cases} \frac{(2q - 2q^3)t - (3q^3 + q^5)t^2}{2 - 2q^2 - 4q^2t} & \text{for } m = 3 \\ \frac{2qt - 3q^m t^2 - q^{m+2}[m-3]_q t^3}{2 - 2q^2 t([m-2]_q + q^{m-3})} & \text{for } 4 \leq m < \infty \\ \frac{qt - q^2 t}{1 - q - q^2 t} & \text{for } m = \infty. \end{cases}$$

Proof. By Theorem 3.1, $|D_R(w)| \leq 2$ for any palindromic $w \in W(m, n)$. Hence we can partition the set of inseparable palindromic elements into those with $|D_R(w)| = 1, 2$ respectively. For notation, let $A_{n, k}$ be the number of inseparable palindromic $w \in W(m, n)$ of length k with $|\mathcal{S}(w)| = n$ and $D_R(w) = 1$. Let $B_{n, k}$ be the number of those same elements with $D_R(w) = 2$. We have $I_{n, k} = A_{n, k} + B_{n, k}$. Consider the polynomials

$$A_n(q) := \frac{1}{n!} \sum_{k \geq 1} A_{n, k} q^k \quad \text{and} \quad B_n(q) := \frac{1}{n!} \sum_{k \geq 1} B_{n, k} q^k.$$

If $n = 1$, then

$$A_1(q) = q \quad \text{and} \quad B_1(q) = 0.$$

If $3 \leq m < \infty$, then for $n = 2$, the inseparable elements have the form $s_1 s_2 s_1 \cdots$ or $s_2 s_1 s_2 \cdots$ where the length is at least 3. There is also a unique longest element $w_0 := \underbrace{s_1 s_2 \cdots}_m$ with $|D_R(w_0)| = 2$. This gives

$$A_2(q) = q^3[m-3]_q \quad \text{and} \quad B_2(q) = \frac{q^m}{2}.$$

For the remainder of the proof, let $w = v_1 \cdots v_{|\mathbf{S}(w)|} \in W(m, n)$ be a complete Grassmannian factorization. We first consider the case when $m = 3$. If w is palindromic and inseparable, then by Theorem 3.1, each v_i is a spiral word as in Theorem 3.1 part 3. In particular, for each even length, there is a unique v_i of up to S_3 permutation symmetry on the generators $\{r, s, t\}$. Moreover, if $|\mathbf{S}(w)| \geq 3$, then $|D_L(w)| = 2$. Thus for all $n \geq 3$, we have $A_n(q) = 0$ and

$$B_n(q) = \left(\frac{2q^2}{1-q^2} \right) B_{n-1}(q) = \frac{q^3}{2} \left(\frac{2q^2}{1-q^2} \right)^{n-2}.$$

Hence

$$\phi_3(q, t) = qt + \frac{q^3 t^2}{2} + \frac{q^5}{1-q^2} \sum_{n \geq 3} \left(\frac{2q^2}{1-q^2} \right)^{n-3} t^n.$$

This proves the first equation in (5).

Now suppose $4 \leq m < \infty$. In this case, if w is palindromic and inseparable, then Theorem 3.1 implies $|\mathbf{S}(v_i)| \leq 2$. Hence each factor v_i has a reduce expression $stst \cdots$ where $t \in D_L(v_1 \cdots v_{i-1})$. In particular, when constructing $w = v_1 \cdots v_{|\mathbf{S}(w)|}$, there are exactly twice as many choices for v_i if $D_L(v_1 \cdots v_{i-1}) = 2$ than if $D_L(v_1 \cdots v_{i-1}) = 1$. This yields that for $n \geq 3$, the polynomials $A_n(q)$ and $B_n(q)$ satisfy the first order recurrence

$$\begin{aligned} A_n(q) &= q^2[m-3]_q (A_{n-1}(q) + 2B_{n-1}(q)) \\ B_n(q) &= q^{m-1} (A_{n-1}(q) + 2B_{n-1}(q)). \end{aligned}$$

This implies that

$$\begin{aligned} \begin{bmatrix} A_n(q) \\ B_n(q) \end{bmatrix} &= \begin{bmatrix} q^2[m-3]_q & 2q^2[m-3]_q \\ q^{m-1} & 2q^{m-1} \end{bmatrix}^{n-2} \begin{bmatrix} A_2(q) \\ B_2(q) \end{bmatrix} \\ &= q^5[m-2]_q (q^2[m-3]_q + 2q^{m-1})^{n-3} \begin{bmatrix} [m-3]_q \\ q^{m-3} \end{bmatrix} \\ &= q^7[m-2]_q ([m-2]_q + q^{m-2})^{n-3} \begin{bmatrix} [m-3]_q \\ q^{m-3} \end{bmatrix}. \end{aligned}$$

Thus

$$\phi_m(q, t) = qt + \left(q^3[m-3]_q + \frac{q^m}{2} \right) t^2 + q^7[m-2]_q^2 \sum_{n \geq 3} ([m-2]_q + q^{m-3})^{n-3} t^n$$

which proves the second equation in (5).

Finally, we compute the exponential generating series for the uniform Coxeter group $W(\infty, n)$ by taking the limit of ϕ_m in the second equation of (5) as $m \rightarrow \infty$. This is equivalent to taking

$$q^m \rightarrow 0 \quad \text{and} \quad [m]_q \rightarrow \frac{1}{1-q}$$

which yields the third equation in (5). $\square$

The following equations are the first few terms in the Taylor expansion of $\Phi_m(q, t)$ for $m = 3, 4, \infty$. These calculations were computed using the `combinat` package for Mupad.

$$\begin{aligned} \Phi_3(q, t) = & 1 + (1 + q)t + (1 + 2q + 2q^2 + q^3) \frac{t^2}{2} \\ & + (1 + 3q + 6q^2 + 9q^3 + 6q^4 + 6q^5 + 6q^7 + O(q^9)) \frac{t^3}{6} \\ & + (1 + 4q + 12q^2 + 30q^3 + 48q^4 + 60q^5 + 54q^6 + O(q^7)) \frac{t^4}{24} + O(t^5) \end{aligned}$$

$$\begin{aligned} \Phi_4(q, t) = & 1 + (1 + q)t + (1 + 2q + 2q^2 + 2q^3 + q^4) \frac{t^2}{2} \\ & + (1 + 3q + 6q^2 + 12q^3 + 15q^4 + 12q^5 + 12q^6 + 6q^7) \frac{t^3}{6} \\ & + (1 + 4q + 12q^2 + 36q^3 + 78q^4 + 120q^5 \\ & + 156q^6 + 168q^7 + 150q^8 + 120q^9 + 48q^{10}) \frac{t^4}{24} + O(t^5) \end{aligned}$$

$$\begin{aligned} \Phi_\infty(q, t) = & 1 + (1 + q)t + (1 + 2q + 2q^2 + 2q^3 + 2q^4 + 2q^5 + O(q^6)) \frac{t^2}{2} \\ & + (1 + 3q + 6q^2 + 12q^3 + 18q^4 + 24q^5 + O(q^6)) \frac{t^3}{6} \\ & + (1 + 4q + 12q^2 + 36q^3 + 84q^4 + 156q^5 + O(q^6)) \frac{t^4}{24} + O(t^5) \end{aligned}$$

By evaluating $\Phi_m(q, t)$ at $q = 1$, we can recover the total number of palindromic elements in $W(m, n)$. By Corollary 3.5, this value is finite only when $4 \leq m < \infty$. We list these values for $4 \leq m \leq 8$ and $1 \leq n \leq 7$ in Figure 3.

4. PROPERTIES OF TRIANGLE GROUPS IN HQ

We discuss a few properties of triangle groups in HQ. The first property is that there are k -palindromic Poincaré polynomials which are not palindromic for large k :

Proposition 4.1. *Let W be the triangle group $(2, b, c)$ with $S = \{r, s, t\}$ such that*

$$(rs)^2 = (rt)^b = (st)^c = e$$

$m \setminus n$	1	2	3	4	5	6	7
4	2	8	67	893	15596	330082	8165963
5	2	10	115	2057	47356	1314292	42584795
6	2	12	175	3893	110436	3768982	150113447
7	2	14	247	6545	219956	8884312	418725119
8	2	16	331	10157	393916	18351562	997538291

FIGURE 3. Number of palindromic elements in $W(m, n)$

where $b, c \geq 3$ and c is finite. Then there exist elements $w \in W$ which are $(c-2)$ -palindromic but not palindromic.

Proof. Consider $w = uv$ where

$$(6) \quad u^{-1} = stst \dots \quad \text{and} \quad v = rtstst \dots$$

with $\ell(u) < c$ and $\ell(v) \leq c$. Calculation of the polynomial $P_w(q)$ reduces to determining the cardinality of the sets

$$M_k := \{w' \leq w \mid \ell(w') = k\}.$$

First we partition

$$M_k = (M_k \cap W_{\{s,t\}}) \sqcup (M_k \cap W \setminus W_{\{s,t\}}).$$

If $w' \in W_{\{s,t\}}$, then w' has the form $sts \dots$ or $tst \dots$. Hence

$$|M_k \cap W_{\{s,t\}}| = \begin{cases} 2 & \text{if } k < \min\{c, \ell(w) - 2\} \\ 1 & \text{if } k = c \text{ or } \ell(w) - 1 \\ 0 & \text{if } k > c. \end{cases}$$

If $w' \in W \setminus W_{\{s,t\}}$, then it is uniquely determined by its parabolic decomposition $w' = u'v'$ where $u' \leq u$, $v' \leq v$ and v' is nontrivial in $W^{\{s,t\}}$. Hence

$$W \setminus W_{\{s,t\}} \simeq [e, u] \times ([r, v] \cap W^{\{s,t\}}).$$

This gives that

$$|M_k \cap W \setminus W_{\{s,t\}}| = \begin{cases} 2k - 1 & \text{if } k \leq \min\{\ell(u), \ell(v)\} \\ 2\ell(v) & \text{if } \ell(v) < k \leq \ell(u) \\ 2\ell(u) & \text{if } \ell(u) < k \leq \ell(v) \\ 2\ell(w) - 2k + 1 & \text{if } k \geq \max\{\ell(u), \ell(v)\} \end{cases}$$

In the case that $\ell(u) = \ell(v) = c - 1$, we have that

$$P_w(q) = [\ell(w) + 1]_q + q^{c+1} + \sum_{k=1}^c 2q^k [\ell(w) - 2k + 1]_q$$

In particular, if we write $P_w(q) = \sum a_i q^i$, then we have that

$$a_i = a_{\ell(w)-i} = 2i + 1$$

for $i \leq c - 3$ and

$$a_{c-2} = 2c - 1, \quad a_c = 2c - 2.$$

Hence $P_w(q)$ is $(c - 2)$ -palindromic but not palindromic. For example, if we take $c = 4$ and $w = uv = (sts)(rts)$, then

$$[e, w] \cap W_{\{s,t\}} = \{e, s, t, st, ts, tst, sts, stst\}$$

and

$$[e, w] \cap W \setminus W_{\{s,t\}} = [e, u] \cdot ([r, v] \cap W^{\{s,t\}}) = \{e, s, t, st, ts, sts\} \cdot \{r, rt, rts\}.$$

In this case, the Poincaré polynomial $P_w(q) = 1 + 3q + 5q^2 + 7q^3 + 6q^4 + 3q^5 + q^6$ is 2-palindromic but not 3-palindromic. $\square$

It is tempting to conjecture that, for the triangle groups $(2, b, c)$ as in Proposition 4.1, all $(c - 1)$ -palindromic elements are palindromic. However for triangle group $(2, 3, 5)$ (Coxeter type H_3 with $c = 5$) there is a unique length 14 element which is 4-palindromic but not palindromic given by $w = tsrtsrtsrtsrtr$.

Theorem 3.1 states that any Grassmannian factorization of a 2-palindromic element $w \in W$ is also a BP-decomposition if W avoids triangles in HQ. This statement is not true for Coxeter groups which contain triangles in HQ.

Proposition 4.2. *Let W be a Coxeter group. Then W avoids all triangle groups in HQ if and only if every Grassmannian factorization $w = uv$ where w is palindromic is a BP-decomposition.*

Proof. By Theorem 3.1, it suffices to show that for triangle groups $(2, b, c)$ as in Proposition 4.1 there are Grassmannian factorizations $w = uv$ of palindromic w which are not BP-decompositions. Consider $w = uv$ as in Equation (6) with $\ell(u) = 2$ and $\ell(v) = c = m_{st}$. It is easy to check that w is palindromic and that $w = uv$ is a Grassmannian factorization with respect to $J = \{s, t\}$ but not a BP-decomposition. $\square$

5. DESCENT SETS OF TRIANGLE AVOIDING GROUPS

In this section, we prove several basic properties of Coxeter groups which avoid triangle groups in HQ. We begin with a lemma that holds for all Coxeter groups:

Lemma 5.1. *Let W be a Coxeter group and $u \in W$. If $s \notin D_L(u)$, then $D_L(su) \setminus \{s\}$ consists of the elements $t \in D_L(u)$ such that u has a reduced factorization starting with a braid $tsts \cdots$ of length $m_{st} - 1$. (If $m_{st} = 2$ then this braid consists of only one element.) In other words,*

$$D_L(su) = \{s\} \cup \{t \in D_L(u) : u = u_0 u_1, u_0 \in W_{\{s,t\}}, u_1 \in W^{\{s,t\}}, \ell(u_0) = m_{st} - 1\}.$$

Proof. Let $J = D_L(su)$. Then by [6], W_J is a finite Coxeter group and su has a reduced factorization beginning with the maximal element w_0 of W_J . If t is an element of $J \setminus \{s\}$, then $m_{st} < \infty$ and w_0 has a reduced decomposition starting with the longest element of $W_{\{s,t\}}$. $\square$

We now consider Coxeter groups which avoid triangle groups in HQ.

Lemma 5.2. *If W is a Coxeter group which avoids all triangle groups in HQ, then the only finite parabolic subgroups of W are products of rank 2 Coxeter groups.*

In other words, if $J \subset S$ is such that W_J is finite, then J can be written as a disjoint union

$$J = \bigsqcup_i J_i,$$

where $|J_i| \leq 2$ for all i , and $m_{st} = 2$ if $s \in J_i$, $t \in J_j$, $i \neq j$.

Proof. Using the classification of finite Coxeter groups, we see that every finite irreducible Coxeter group of rank ≥ 3 contains a triangle group in HQ. $\square$

If $J = D_L(w)$, then W_J is a finite Coxeter group. In particular, Lemma 5.2 applies to the parabolic subgroups generated by descent sets of HQ-avoiding Coxeter groups. The following lemma is the main result of this section.

Lemma 5.3. *Let (W, S) be a Coxeter group which avoids triangle groups in HQ. Let $r, s \in S$ such that $3 \leq m_{rs} \leq \infty$, and suppose u is an element of W such that $(rs)u$ is a reduced factorization. Then*

$$D_L(rsu) \setminus \{r, s\} = \{t \in D_L(u) : m_{rt} = m_{st} = 2\}.$$

Proof. The proposition is obviously true if $u = e$. We proceed by induction on the length of u . Let $J = D_L(su)$, and write $J = \bigsqcup J_i$ as in Lemma 5.2. We can further assume that if $J_i = \{x, y\}$, then $m_{xy} \geq 3$, and that $s \in J_0$.

Now if $t \in D_L(rsu) \setminus \{r, s\}$, then by Lemma 5.1 we must have $m_{rt} < \infty$ and rsu must have a reduced decomposition starting the longest element in $W_{\{r,t\}}$. If $t \notin J_0$ then $m_{st} = 2$. Since W avoids triangle groups in HQ, we have that $m_{rt} = 2$ as well.

This leaves the possibility that $t \in J_0$, in which case $m_{st} \geq 3$. Once again, since W avoids triangle groups in HQ, we conclude that $m_{rt} \geq 3$. Thus rsu has a reduced factorization $rsu = (rtr)u'$, where $\ell(u') = \ell(u) - 1$. Now $tru' = su$, so $s \in D_L(tru')$. But by induction, this implies that $m_{ts} = m_{rs} = 2$, which is a contradiction. Hence $t \notin J_0$. $\square$

6. PROOF OF THEOREM 3.1

We now prove Theorem 3.1. The following assumptions are fixed for the remainder of the section. Let W be a Coxeter group that avoids all triangle groups in HQ. Let $w \in W$ be 2-palindromic with a Grassmannian factorization $w = uv$ with respect to $J = S(u)$. By Lemma 2.6 we have that

$$(7) \quad |u \cdot D(v) \cap D(w)| = 1.$$

This implies

$$(8) \quad |W^J \cap D(v)| = 1,$$

and in particular, $|D_R(v)| = 1$. Let $z \in D_R(v)$ denote this unique simple reflection. The element vz is the unique element in $W^J \cap D(v)$.

We divide the proof into three steps. The first step is to prove that $S(v)$ has at most three elements. Second, we prove the characterization of v when $S(v)$ has

exactly three elements. For the last step, we show that $w = uv$ is BP-decomposition. We begin with the following technical lemma.

Lemma 6.1. *Let $s_1, \dots, s_k$ be the longest sequence of distinct simple reflections such that v has a reduced decomposition*

$$v = s_1 \cdots s_k v',$$

and for all $j < k$, $m_{s_j s_{j+1}} \geq 3$. For any $1 \leq j \leq k$, define the set $I_j := \{s_1, \dots, s_j\}$. Then:

- (1) $s_j \cdots s_k v' \in W^{I_{j-1}}$ for all $j \leq k$, and
- (2) $S(v') \subseteq \{s_1, \dots, s_k\}$.

Proof. Clearly the lemma is true if $\ell(v) = 1$ and hence we assume that $\ell(v) \geq 2$. Observe that $k \geq 2$, otherwise $v \notin W^J$. For any $j \leq k$, let

$$v = v_j v'_j$$

be a parabolic decomposition with respect to W_{I_j} . It is easy to see that $v_1 = s_1$ and hence $v'_1 = s_2 \cdots s_k v' \in W^{I_1}$.

Now let $j \geq 2$ and suppose that $\ell(v_j) > j$. Then there exists $s \in D_R(v_j)$ such that

$$S(v_j s) = S(v_j).$$

By Lemma 5.3, we have that the left descent sets

$$D_L(v_j s v'_j) = D_L(v_j s) \cup \{t \in D_L(v'_j) \mid m_{ts_i} = 2 \text{ for } i \leq j\}$$

and

$$D_L(v) = D_L(v_j v'_j) = D_L(v_j) \cup \{t \in D_L(v'_j) \mid m_{ts_i} = 2 \text{ for } i \leq j\}.$$

Since $S(v_j s) = S(v_j)$, the descent sets above are equal. Hence $v_j s v'_j \in W^J \cap D(v)$. If $j < k$, then $v'_j \neq e$ and consequently $v_j s v'_j \neq v z$, contradicting Equation (8). Thus $\ell(v_j) = j$ which implies that $v_j = s_1 \cdots s_j$ and $v'_j = s_{j+1} \cdots s_k v' \in W^{I_j}$. This proves part (1) of the lemma.

For part (2), suppose that $S(v') \not\subseteq \{s_1, \dots, s_k\}$. Then $|S(v)| > k$ and $v'_k \neq e$. We get that $v_k = s_1 \cdots s_k$ and

$$D_L(v'_k) \cap \{s_1, \dots, s_k\} = \emptyset$$

since $v'_k \in W^{I_k}$. By the maximality of k , we have that $m_{s_k, r} = 2$ for all $r \in D_L(v'_k)$. We claim that

$$D_L(s_1 \cdots s_{k-1} v') = D_L(v).$$

Indeed, if $k \geq 3$, this follows from Lemma 5.3. Otherwise, if $k = 2$, then

$$m_{s_1 s_2} = m_{s_1 r} = \infty$$

since W avoids all triangle groups in HQ . This proves the claim when $k = 2$. In either case we have that

$$s_1 \cdots s_{k-1} v' \in W^J \cap D(v)$$

which contradicts Equation (8). Therefore $S(v') \subseteq \{s_1, \dots, s_k\}$. $\square$

The following proposition completes the first step of in the proof of Theorem 3.1.

Proposition 6.2. *We have that $|\mathbf{S}(v)| \leq 3$. Furthermore, if $|\mathbf{S}(v)| = 3$, then $\mathbf{S}(v)$ generates a triangle group (a, b, c) with $a, b, c \geq 3$.*

Proof. Suppose $|\mathbf{S}(v)| \geq 4$ and let $v = s_1 \cdots s_k v'$ as in Lemma 6.1. We first show by induction on j that

- (1) $D_L(s_j \cdots s_k v') = \{s_j\}$
- (2) $m_{s_i s_j} = 2$ for $i \in \{1, \dots, j-2\}$.

Indeed, part (1) is trivial for $j = k$. Suppose part (1) is true for some $j \leq k$. Now by Lemma 6.1, $s_1 \cdots s_{j-2} s_j \cdots s_k v'$ is reduced, and therefore is not an element of W^J . So by Lemma 5.3, we have that $s_j \in D_L(s_1 \cdots s_{j-2} s_j \cdots s_k v')$. Moreover, if $j \geq 4$, then $m_{s_i s_j} = 2$ for all $1 \leq i \leq j-2$. If $j = 3$, then $s_1 s_3 \cdots s_k v'$ has a reduced expression beginning with a braid $s_1 s_3 s_1 \cdots$ of length $m_{s_1 s_3} < \infty$. Since $s_1 \notin D_L(s_4 \cdots s_k v')$, we conclude that $m_{s_1 s_3} = 2$. Hence part (2) holds for j .

Now suppose part (2) holds for all $j > j_0$. Since $|D_L(v)| = 1$, we have that $s_j \notin D_L(s_{j_0} \cdots s_k v')$ for any $j > j_0$. Thus part (1) holds for j_0 . Hence (1) and (2) hold for all j .

Now part (2), combined with the HQ-avoiding condition, implies that

$$m_{s_i s_j} = \begin{cases} \infty & |i - j| = 1 \\ 2 & |i - j| \geq 2 \end{cases}.$$

In other words, if $|\mathbf{S}(v)| \geq 4$ then $W_{\mathbf{S}(v)}$ is defined entirely by commuting relations. We show that this hypothesis implies that $|\mathbf{S}(v)| \leq 2$. Indeed, suppose $|\mathbf{S}(v)| \geq 3$, and let $u = u_1 u_0$, where $u_1 \in {}^{\mathbf{S}(v)}W$ and $u_0 \in W_{\mathbf{S}(v)}$. Here the set ${}^{\mathbf{S}(v)}W$ denotes the minimal length representatives of the left cosets $W/W_{\mathbf{S}(v)}$. By Equation (7), the product $u_0 s_2 \cdots s_k v'$ must not be reduced. We conclude that $D_R(u_0) \cap D_L(s_2 \cdots s_k v') = \{s_2\}$ since for any $s_i, s_j \in \mathbf{S}(u_0) \cup \mathbf{S}(v) = \mathbf{S}(v)$ we have $m_{s_i s_j} = 2$ or ∞ . Moreover, since $D_L(s_1 s_3 \cdots s_k v') = \{s_1, s_3\}$, the same argument shows that $s_3 \in D_R(u_0)$. But now we have $\{s_2, s_3\} \subseteq D_R(u_0)$ which implies that the m_{s_2, s_3} is finite. This contradicts the fact that $m_{s_2, s_3} = \infty$. Hence, $|\mathbf{S}(v)| \leq 3$.

Finally, if $|\mathbf{S}(v)| = 3$, then by Lemma 6.1, $m_{s_1 s_2}, m_{s_2 s_3} \geq 3$. If $m_{s_1 s_3} = 2$, then the HQ-avoiding condition implies $m_{s_1 s_2} = m_{s_2 s_3} = \infty$. We can now apply the previous argument as above show that $\{s_2, s_3\} \subseteq D_R(u_0)$ and hence m_{s_2, s_3} is finite. Thus we must have $m_{s_1 s_3} \geq 3$. This completes the proof. $\square$

For the next step in the proof of Theorem 3.1, suppose that $|\mathbf{S}(v)| = 3$ with $\mathbf{S}(v) = \{r, s, t\}$. By Proposition 6.2, we have $m_{rs}, m_{rt}, m_{st} \geq 3$. Consider the reduced factorization $v = xy$ where

$$x^{-1} := tsrtstr \cdots$$

is the largest spiral word prefix of v . In other words, we can write

$$(9) \quad v = xy = (\cdots rstrst) \cdot y.$$

Define $x' := xtst$. It is easy to see that $\ell(x') = \ell(x) - 1$ and that x' equals x with the second to last reflection s removed. For any $0 \leq k \leq \ell(x')$ define a length k suffix x'_k

of x' by

$$x'_k := \underbrace{\cdots rstrt}_k$$

Lemma 6.3. *For any $0 \leq k \leq \ell(x')$, the following are true:*

- (1) *The product $x'_k y$ is a reduced factorization.*
- (2) *If k is even, then $|D_L(x'_k y)| = 1$. If k is odd, then $|D_L(x'_k y)| \leq 2$.*
- (3) *If $|D_L(x'_k y)| = 2$ and $k \geq 5$, then $|D_L(x'_{k-2} y)| = 2$.*

Proof. If $k = 0$, then $r, t \notin D_L(y)$ since x is a maximal length spiral word. This implies that $D_L(y) = \{s\}$. If $k = 1$, then ty is reduced and $D_L(ty) \subseteq \{s, t\}$. Moreover, $rtty$ is reduced and by Lemma 5.3, we have that $D_L(rty) = \{r\}$ since $r \notin D_L(y)$. This proves the lemma for $k \leq 2$.

We proceed with the proof by induction on k . Suppose $k \geq 3$. Without loss of generality, we can assume $r \in D_L(x'_k)$, so that s is the first element of x'_{k-1} . We first consider the case where k is odd. Then by the inductive assumption, we have that $D_L(x'_{k-1} y) = \{s\}$. Hence

$$x'_k y = r x'_{k-1} y$$

is reduced and $D_L(x'_k y) \subseteq \{r, s\}$. If k is even, then s and t are the first two elements of x'_{k-1} ; in particular, r is not one of the first two elements. Therefore

$$D_L(x'_{k-2} y) = \{t\} \quad \text{and} \quad D_L(x'_{k-1} y) \subseteq \{s, t\}.$$

So $x'_k y$ is reduced and $D_L(x'_k y) = \{r\}$. This proves parts (1) and (2) of the lemma.

To prove part (3), suppose that $k \geq 5$ is odd with

$$t \in D_L(x'_{k-2} y) \subseteq \{t, r\} \quad \text{and} \quad r \in D_L(x'_k y) \subseteq \{r, s\}.$$

If $|D_L(x'_k y)| = 2$, then $r \in D_L(x'_{k-2} y)$ since $3 \leq m_{rs} < \infty$. Hence $|D_L(x'_{k-2} y)| = 2$. $\square$

One immediate consequence of Lemma 6.3 is that $x'y$ is a reduced factorization and that if $\ell(x')$ is even, then $x'y \in W^J \cap D(v)$ which is a contradiction to Equation (8). Hence $\ell(x')$ is odd (i.e. $\ell(x)$ is even). The following lemma is a preliminary characterization of v .

Lemma 6.4. *The spiral word x satisfies one of the following conditions:*

- (1) $m_{rt} = 3$ and $\ell(x) = 4$.
- (2) $m_{rt} = m_{rs} = 3$ and $\ell(x) = 6$.
- (3) $m_{rt} = m_{rs} = m_{st} = 3$ and $\ell(x) \geq 8$.

Proof. Since $x'y$ is reduced, we have that $x'y \notin W^J$ and $\ell(x') \geq 3$. Furthermore, by Lemma 6.3 part (3), $|D_L(x'_k y)| = 2$ for all $k \geq 3$. In particular the following statements are true:

- For $k = 3$, we have that $|D_L(trty)| = 2$ if and only if $m_{rt} = 3$.
- For $k = 5$, we have that $|D_L(rstrty)| = 2$ if and only if $m_{rt} = m_{rs} = 3$.
- For $k = 7$, we have that $|D_L(strstrty)| = 2$ if and only if $m_{rt} = m_{rs} = m_{st} = 3$.

This completes the proof. $\square$

Now we consider the reduced factorization

$$(10) \quad v = xy = (\cdots rstrst) \cdot y = \overbrace{(\cdots rstr)}^x \underbrace{(st \ st \ \cdots)}_{\text{length } k} \cdot \bar{y},$$

where k is the length of the longest possible prefix of sty the form $stst \cdots$.

Lemma 6.5. *With v as Equation (10), the following are true:*

- (1) $\bar{y} = e$.
- (2) $k = m_{st} - 1$

Proof. Suppose that $\bar{y} \neq e$. Then $D_L(\bar{y}) = \{r\}$ by the maximality of k . If $k = 2$, then x is not a maximal length spiral, and hence $\bar{y} = e$. Now assume that $k \geq 3$ and let $v = \bar{x}\bar{z}\bar{y}$ be the reduced factorization given in (10) where $\bar{z} \in W_{\{s,t\}}$ is of length k . Without loss of generality, let $t \in D_R(\bar{z})$ and define $\bar{z}' := \bar{z}t$. Since $k \geq 3$, we have $\ell(\bar{z}') \geq 2$ and thus $\bar{x}\bar{z}'\bar{y}$ is a reduced factorization. Likewise, since $\ell(\bar{z}') \geq 2$ and $D_R(\bar{x}) = \{r\}$, we have that $\bar{x}\bar{z}'\bar{y} \in W^J$ and hence $\bar{x}\bar{z}'\bar{y} \in W^J \cap D(v)$. But this contradicts Equation (8). Therefore $\bar{y} = e$ and part (1) of the lemma is proved.

Since $\ell(x)$ is even, we have that $k < m_{st}$, otherwise $v \notin W^J$. This completes the proof in the case of x as in Lemma 6.4 part (3). Now suppose that $k \leq m_{st} - 2$. If x satisfies the condition in Lemma 6.4 part (1), then $J = S(w) \setminus \{t\}$ and we can write

$$v = tr \underbrace{stst \cdots}_{\text{length } k} = tr\bar{z}.$$

But then $t\bar{z} \in W^J \cap D(v)$ which contradicts Equation (8). If x satisfies the condition in Lemma 6.4 part (2), then $J = S(w) \setminus \{r\}$ and

$$v = rstr \underbrace{stst \cdots}_{\text{length } k} = rstr\bar{z}.$$

But then $rstr\bar{z} \in W^J \cap D(v)$ which also contradicts Equation (8). Hence $k > m_{st} - 2$ and part (2) of the lemma is proved. $\square$

It is easy to see that Lemmas 6.3, 6.4 and 6.5 prove the characterization v when $|S(v)| = 3$ in Theorem 3.1.

The final step in the proof is to show that $w = uv$ is a BP-decomposition. In this step, we do not assume that $|S(v)| = 3$.

Lemma 6.6. *For any $s_0 \in S(v) \cap J$, there exists $v'' \in W^J$ of length $\ell(v'') = \ell(v) - 2$ such that $s_0 v'' \in D(v)$.*

Proof. If $|S(v)| \leq 2$, then the lemma is obvious. If $|S(v)| = 3$, then we can write $v = xy$ as in Equation (9) with the notational change that

$$x = rstrst \cdots.$$

In other words, we let r, s, t denote the first three simple reflections appearing in x , rather than the last three. We want to find v'' for $s_0 \in S(v) \cap J = \{s, t\}$. Note that with the change in notation, we have that $m_{rs} = 3$. Recall the definition of x' given

after Equation (9). By Lemma 6.3 part (1) we have that $x'y$ is reduced and hence $x'y \in D(v) \cap W^{\{r,s\}}$. Thus we have a reduced factorization

$$x'y = (srs)y'$$

for some y' . For $s_0 = s$, we set $v'' = rsy'$. Then $v'' \in W^J$ since $D_L(v'') = \{r\}$.

We now find a v'' for $s_0 = t$. Consider the reduced factorization

$$v = (rs)(ty'').$$

Clearly $r \notin D_L(ty'')$, otherwise $v \notin W^J$. Hence $rtty'' \in D(v)$ and $rtty'' \notin W^J$. This implies that $t \in D_L(rtty'')$ and we can write a reduced factorization

$$rtty'' = (trt)y'''$$

for some y''' . We set $v'' = rtty'''$ for $s_0 = t$. Since $D_L(v'') = \{r\}$, we get that $v'' \in W^J$. This completes the proof. $\square$

If $s_0 \in S(v) \cap J$ and $v'' \in W^J$, such that $s_0v'' \in D(v)$, then $s_0 \in D_R(u)$. Otherwise $us_0v'' \in u \cdot D(v) \cap D(w)$ which contradicts Equation (7). Applying Lemma 2.2, we get that $w = uv$ is a BP-decomposition. This completes the proof of Theorem 3.1.

Acknowledgements. The second author would like to thank the University of British Columbia for its hospitality. The authors thank Sara Billey for her useful emails and Jim Carrell and Alex Woo for some helpful discussions. The authors would also like to thank the referees for their suggestions on improving the manuscript. The second author was partially supported by NSF grant DMS-1007255.

REFERENCES

- [1] Alan F. Beardon. *The geometry of discrete groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1983.
- [2] Sara Billey and Andrew Crites. Rational smoothness and affine Schubert varieties of type A . In *23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011)*, Discrete Math. Theor. Comput. Sci. Proc., AO, pages 171–181. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2011.
- [3] Sara Billey and V. Lakshmibai. *Singular loci of Schubert varieties*, volume 182 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2000.
- [4] Sara Billey and Alexander Postnikov. Smoothness of Schubert varieties via patterns in root subsystems. *Adv. in Appl. Math.*, 34(3):447–466, 2005.
- [5] Sara C. Billey. Pattern avoidance and rational smoothness of Schubert varieties. *Adv. Math.*, 139(1):141–156, 1998.
- [6] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [7] Anders Björner and Torsten Ekedahl. On the shape of Bruhat intervals. *Ann. of Math. (2)*, 170(2):799–817, 2009.
- [8] Miklós Bóna. The permutation classes equinumerous to the smooth class. *Electron. J. Combin.*, 5:Research Paper 31, 12 pp. (electronic), 1998.
- [9] James B. Carrell. The Bruhat graph of a Coxeter group, a conjecture of Deodhar, and rational smoothness of Schubert varieties. In *Algebraic groups and their generalizations: classical methods (University Park, PA, 1991)*, volume 56 of *Proc. Sympos. Pure Math.*, pages 53–61. Amer. Math. Soc., Providence, RI, 1994.
- [10] Vesselin Gasharov. Factoring the Poincaré polynomials for the Bruhat order on S_n . *J. Combin. Theory Ser. A*, 83(1):159–164, 1998.

- [11] David Kazhdan and George Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53(2):165–184, 1979.
- [12] David Kazhdan and George Lusztig. Schubert varieties and Poincaré duality. In *Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, Proc. Sympos. Pure Math., XXXVI, pages 185–203. Amer. Math. Soc., Providence, R.I., 1980.
- [13] V. Lakshmibai and B. Sandhya. Criterion for smoothness of Schubert varieties in $Sl(n)/B$. *Proc. Indian Acad. Sci. Math. Sci.*, 100(1):45–52, 1990.
- [14] Stephen A. Mitchell. A filtration of the loops on $SU(N)$ by Schubert varieties. *Math. Z.*, 193(3):347–362, 1986.
- [15] Suho Oh and Hwanchul Yoo. Bruhat order, rationally smooth Schubert varieties, and hyperplane arrangements. *DMTCS Proceedings*, FPSAC, 2010.
- [16] Zvezdelina E. Stankova. Forbidden subsequences. *Discrete Math.*, 132(1-3):291–316, 1994.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA

E-mail address: erichmond@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA

E-mail address: slofstra@math.berkeley.edu